

# Does rotation of B stars depend on metallicity? preliminary results from GIRAFFE spectra

Frédéric Royer<sup>1</sup>, Pierre North<sup>2</sup>, Claudio Melo<sup>3</sup>, Jean-Claude Mermilliod<sup>2</sup>,  
Eva K. Grebel<sup>4</sup>, Jose Renan de Medeiros<sup>5</sup>, and André Maeder<sup>6</sup>

<sup>1</sup> GEPI / CNRS UMR 8111, Observatoire de Paris, F-92195 MEUDON cedex, France

<sup>2</sup> Laboratoire d'Astrophysique, Ecole Polytechnique Fédérale de Lausanne (EPFL),  
Observatoire, CH-1290 Sauverny, Switzerland

<sup>3</sup> ESO, Alonso de Cordova 3107, Casilla 19001, Santiago 19, Chile

<sup>4</sup> Astronomisches Institut, Universität Basel, Venusstr. 7, CH-4102 Binningen,  
Switzerland

<sup>5</sup> Universidade Federal do Rio Grande do Norte, 59072-970 Natal, R.N., Brazil

<sup>6</sup> Observatoire de Genève, CH-1290 Sauverny, Switzerland


**Abstract.** We show the  $v \sin i$  distribution of main sequence B stars in sites of various metallicities, in the absolute magnitude range  $-3.34 < M_V < -2.17$ . These include Galactic stars in the field measured by [1], members of the  $\eta$  &  $\chi$  Per open clusters measured by [6], and five fields in the SMC and LMC measured at ESO Paranal with the FLAMES-GIRAFFE spectrograph, within the Geneva-Lausanne guaranteed time. Following the suggestion by [5], we do find a higher rate of rapid rotators in the Magellanic Clouds than in the Galaxy, but the  $v \sin i$  distribution is the same in the LMC and in the SMC in spite of their very different metallicities.

## 1 Introduction, results and conclusion

This work aims at testing the suggestion of [5] that stellar rotation is faster at lower metallicity by direct measurements, especially in the LMC and SMC, on stars with  $-3.34 < M_V < -2.17$ , i.e. spectral types B0-B6 or masses from  $\sim 6.7$  to  $14 M_\odot$ . This work is complementary to that of [4], which deals with slightly more massive stars. The results are shown on Fig. 1 and commented in the caption. There is an excess of slow rotators in the Galaxy relative to the MCs, but the  $v \sin i$  distributions of the LMC and the SMC are surprisingly similar.

## References

1. H.A. Abt, H. Levato, M. Grosso: ApJ **573**, 359 (2002)
2. N. Cramer: New Astronomy Reviews **43**, 343 (1999)
3. D. Erspamer, P. North: A&A **383**, 227 (2002)
4. S.C. Keller: Pub. Astron. Soc. Australia **21**, 310 (2004)
5. A. Maeder, E. Grebel, J.-C. Mermilliod: A&A **346**, 459 (1999)
6. P. North, F. Royer, C. Melo et al.: 'New homogeneous  $v \sin i$  determinations for B stars in galactic open clusters'. In: *Stellar Rotation, IAU Symp. 215*, ed. by A. Maeder and P. Eenens, ASP, p. 69
7. A. Slettebak, G.W. Collins, T.D. Parkinson et al.: ApJS **29**, 137 (1975)
8. J. Southworth, P.F.L. Maxted, B. Smalley: MNRAS **349**, 547 (2004)



**Fig. 1.** Cumulative  $v \sin i$  distributions for Galactic stars in the field (dotted line), for members of the h &  $\chi$  Per clusters (dash-dot) and for stars in the LMC (short dash) and in the SMC (solid line). The GIRAFFE spectrograph, attached to the UT2 telescope (VLT) and used in the L2 setup ( $R = 6400$ ,  $\lambda_c = 4272 \text{ \AA}$ ), was used on 3 fields in the LMC (centered on  $[\alpha_{J2000} = 05:31:40, \delta_{J2000} = -66:59:48]$ , on  $[05:30:40, -67:17:12]$  and on  $[05:03:48, -69:00:36]$ ) and 2 fields in the SMC (centered on  $[00:56:12, -72:29:00]$  and on  $[00:49:26, -73:12:07]$ ). We fitted synthetic spectra to observed ones in the range  $4460 - 4490 \text{ \AA}$  with the technique described by [3] using an average  $T_{\text{eff}} - M_V$  relation for the main sequence and assuming  $\log g = 4.0$ . The resulting  $v \sin i$  values were then transformed to the scale of [7]. For the Galaxy, we defined the  $v \sin i$  distribution using 1) the measurements made in the h &  $\chi$  Per clusters by [6] and 2) the large sample of [1] of bright field B stars, Geneva photometry being used to determine  $M_V$  through the calibration of [2]. The SB2 systems were eliminated from this sample, which, although magnitude-limited, does not significantly differ from a volume limited one. The results are summarized in this Figure. Surprisingly, the overall  $v \sin i$  distribution is almost exactly the same in the SMC (mean metallicity  $Z \sim 0.008$ ) and in the LMC ( $Z \sim 0.004$ ). There is only a marginal difference between h &  $\chi$  Per ( $Z(\text{h Per}) \sim 0.01$  according to [8]) and the MCs, but a very significant one ( $P < 0.1 \%$ ) between the Galactic field ( $Z \sim Z_{\odot} = 0.018$ ) and the MC fields. Thus, either the metallicity effect saturates for  $Z < Z(\text{LMC}) \sim 0.008$ , or another cause affects rotational velocities, e.g. different rates and orbital parameters of SB1 binaries (not excluded from the samples), through tidal effects.